

راهنمای آموزشی Clinical Key مغری گلمغانی

کارشناس ارشد علم اطلاعات و دانش شناسی گرایش مدیریت اطلاعات

پایگاه اطلاعاتی Clinical Key

از سال ۲۰۱۲ توسط شرکت Elsevier طراحی شد. این پایگاه جایگزین MD Consult شده است.

ویژگی های Clinical Key


جامعیت: سوالات شما را از طریق مجموعه منابع پزشکی منتشر شده در موضوعات تخصصی پزشکی وجراحی پاسخ می دهد


در برگیرنده خلاصه مقاله MEDLINE، که هر روز از کتابخانه ملی پزشکی (NLM) بازیابی و به صورت یکپارچه به نتایج جستجو Clinical Key اضافه می شود.

قابلیت اعتماد: دارای محتوای معتبر و قابل اطمینان در حوزه پزشکی و جراحی می باشد.

سرعت در پاسخگویی: مرتبط ترین پاسخ ها را در کوتاهترین زمان ارائه میدهد.

جهت ثبت نام در پایگاه گزینه Register را کلیک نمایید. در صفحه گشوده شده اطلاعات را تکمیل کنید. سپس ایمیل ارسالی به پست الکترونیکی خود با تارید نمایید





برای استفاده در دفعات دیگر بایستی نام کاربری و رمزعبور را وارد نموده گزینه Login را کلیک نمایید



ادامه















clinicalkey@elsevier.com ⊘ to Me

Dear Soghra Golmaghani,

You have been sent your username and a link to reset your password by

We found the following username(s) that match your email addge

Username: s.golmaghani@arums.ac.ir Reset password

Click the reset password link for each username above you would like to reset.

Reset password links will expire after 30 minutes.

Thank you, The ClinicalKey Help Desk

Copyright © 2016 Elsevier B.V. | Privacy Policy

Elsevier B.V., Radarweg 29, 1043NX Amsterdam, The Netherlands, Reg. no. 33156677, VAT no. NL 005033019B0

وارد ایمیل خود شوید سپس گزینه Reset Password را کلیک نمایید.

you have completed the Forgotten Password form

در صفحه گشوده شده رمز جدید برای خود تعریف نمایید.

Dogot magazir

Reset password

To complete your password reset, enter a new

word below.


New password


Confirm password

Submit >

https://www.clinicalkey.com

پروفایل اشخاص






نوار جستجو منوی کشویی ALL شامل:

- Books: کتابها
- ا Journals : مجلات
- Clinical Trials : مطالعات بالینی، کار آز مایی بالینی
- Drug Monographs: اطلاعات دارویی، امکان جستجوی بیماری و نمایش دارو های مرتبط با آن و مشاهده اطلاعات دارویی
 - Guidelines > دستور العمل ها
 - ▶ Patient Education: آموزش بيمار
 - < First Consult/Clinical overviews : تشخیص افتراقی براساس پزشکی مبتنی بر شواهد که کاربرد بالینی دارد.
 - ✓ Multimedia: شامل تصاوير و فيلم ها
 - < Procedures Consult: آموزش و راهنمایی گام به گام روند بیش از 350 تکنیک جراحی درمانی و تشخیصی به صورت فیلم و انیمیشین

کلیدواژه مورد نظر را وارد نمایید(در صورتی که املای واژه غلط باشد. این پایگاه املای صحیح را ارائه می دهد)با استفاده از فیلتر سمت چپ، نتایج را محدود نمایید






allergy drugs



متن كامل



□ 43 results

[+] Rate Results

■ MEDLINE®

The diagnosis and management of antibiotic allergy in children: Systematic review to inform a contemporary...

Archives of disease in childhood.

Marrs, Tom; Fox, Adam T... Show all. Published June 1, 2015.

☐ FULL TEXT ARTICLE

Harmonisation in study design and outcomes in paediatric antibiotic clinical trials: a systematic review

Lancet Infectious Diseases, The.

Folgori, Laura, MD; Bielicki, Julia, MD... Show all. Published September 1, 2016. Volume 16, e178-e189. © 2016.

□ FULL TEXT ARTICLE

Endometriosis and type 1 allergies/immediate type hypersensitivity: a systematic review №

European Journal of Obstetrics & Gynecology and Reproductive Biology.

Bungum, Helle Folge; Vestergaard, Christian; Knudsen, Ulla Breth. Published August 1, 2014. Volume 179. Pages 209-215. © 2014.

■ MEDLINE®

A systematic review: can one prescribe carbapenems to patients with IgE-mediated allergy to penicillins or...

Clinical infectious diseases: an official publication of the Infectious Diseases Society of America. Kula, Brittany; Djordjevic, Gordana; Robinson, Joan L. Published October 15, 2014.

☐ FULL TEXT ARTICLE

Efficacy of photodynamic therapy versus antibiotics as an adjunct to scaling and root planing in the...

Drug Allergy Disease Overview View Full Topic Conn's Current Therapy 2017 · Bope, Edward T., MD; Kellerman, Rick D., MD Epidemiology Drug hypersensitivity reactions are one of the many different types of adverse drug reactions (ADRs). ADRs are common, yet the more severe reactions have been estimated to cause 3% to 6% of hospital admissions. More than 100,000 deaths annually are caused by serious ADRs, making these reactions one of the leading cause of death in the United States. Early detection of all ADRs can potentially improve patient outcomes and lower health care costs. More

Was this helpful? Yes or No

Clinical Manifestations

costeroids

Conclusion

Pathophysiology

سرفصل مطالب جهت دسترسی به بخش های مختلف مقاله کلیک نمایید

Introduction

Methods

Search strategy and selection criteria

Results

Study selection and description

Quality assessment of included studies

Patient inclusion criteria

Patient exclusion criteria

Primary and secondary endpoints

Timing of endpoints

Similarities with, and divergence from, adult EMA and FDA guidelines

Discussion

References

پرینت، ایمیل، ذخیره، CME (آموزش مداوم پزشکی) فقط برای کتابها، First Consult و مقالات تمام متن در دسترس می باشد

FULL TEXT ARTICLE

Harmonisation in study design and outcomes in paediatric antibiotic clinical trials: a systematic review \$\infty\$ \$\frac{1}{2}\$

Laura Folgori MD, Julia Bielicki MD, Beatriz Ruiz MD, Mark A Turner MD, John S Bradley Prof, Daniel K Benjamin Prof, Theoklis E Zaoutis Prof, Irja Lutsar Prof, Carlo Giaquinto Prof, Paolo Rossi Prof and Mike Sharland Prof Lancet Infectious Diseases, The, 2016-09-01, Volume 16, Issue 9, Pages e178-e189, Copyright © 2016 Elsevier Ltd

Summary

There is no global consensus on the conduct of clinical trials in children and neonates with complicated clinical infection syndromes. No comprehensive regulatory guidance exists for the design of antibiotic clinical trials in neonates and children. We did a systematic review of antibiotic clinical trials in complicated clinical infection syndromes (including bloodstream infections and community-acquired pneumonia) in children and neonates (0–18 years) to assess whether standardised European Medicines Agency (EMA) and US Food and Drug Administration (FDA) guidance for adults was used in paediatrics, and whether paediatric clinical trials applied consistent definitions for eligibility and outcomes. We searched MEDLINE, Cochrane CENTRAL databases, and ClinicalTrials.gov between Jan 1, 2000, and Nov 18, 2015. 82 individual studies met our inclusion criteria. The published studies reported on an average of 66% of CONSORT items. Study design, inclusion and exclusion criteria, and endpoints varied substantially across included studies. The comparison between paediatric clinical trials and adult EMA and FDA guidance highlighte that regulatory definitions are only variably applicable and used at present. Absence of consensus for paediatric antibiotic clinical trials is a major barrier to harmonisation in research and translation into clinical practice. To improve comparison of therapies and strategies, international collaboration among all relevant stakeholders leading to harmonised case definitions and outcome measures is needed.



THE LANCET Infectious Diseases

Miles and Artifica Ar

Lancet Infectious Diseases, The Volume 16, Issue 9

Copyright © 2016 Elsevier Ltd

New Issue Alerts: Subscribe

Get rights and content 😝


نمونه مقاله بازيابي شده

مرور ليست كتابها

Books Journals Drug Monographs Guidelines Patient Education First Consult/Clinical Overviews Multimed

ليست تخصص ها

Browse Books




Featured Books



Abeloff's Clinical Oncology

روی عنوان کتاب کلیک نمایید صفحه فهرست کتاب باز می شود. اگر روی هر یک از عناوین فهرست کلیک کنید. صفحه مربوطه گشوده می شود



11. Viruses and Human Cancer

Oral Complications

83. Bladder Cancer

underscore the complexity of carcinogenesis and have profoundly influenced the design of strategies for both

محدود کردن براساس تخصص باکس جستجوی مجلات براساس عنوان

مرور ليست مجلات



با استفاده از این گزینه (افزودن Feed آن به Reader خود) می توانید اتوماتیک وار فهرست مقالات جدید را دریافت نمایید

صفحه خانگی مجله JACC

Cardiology)

New Issue Alerts: Subscribe

Search this Journal

در این باکس می توانید کلیدواژه خود را فقط در این مجله جستجو نمایید

Q



Current Issue

2015-6-23, Volume 65, Issue 24

Journal Volumes:

- > Articles in Press
- > Volume 70 (2017)
- > Volume 69 (2017)
 - > Volume 68 (2016)
 - > Volume 67 (2016)
 - > Volume 66 (2015)



Multimedia


Journals **Drug Monographs** Guidelines Patient Education First Consult/Clinical Overviews Books


Multimedia

تصوير


محدود نمودن یافته ها براساس تخصص


محدود نمودن یافته ها براساس تصاوير يا ويدئو





• مشاهده اندازه کامل تصویر • مشاهده تصویر در منبع مربوطه











Fig. 65-1 (Adapted from http://tpis.upmc.com/tpislibrary/HB/H00034f.html .) Adenovirus hepatitis

"smudgy" appearance of the nucleus (arrows), the peripheralized chromatin pattern, and the lack of cytoplasmic inclusions that help differentiate these inclusions from those seen with CMV.

showing characteristic intranuclear inclusion bodies. A high magnification of the adenoviral lesion shows the light microscopic appearance of the typical adenoviral nuclear inclusions. The

ایجاد PowerPoints





Viral Hepatitis in Childre

Zakim and Boyer's Hepatology.


قبل از ایجاد پاورپوینت حتما در پایگاه Clinical Key ثبت نام نمایید.

- 1. تصویر مورد نظر را انتخاب سپس روی آیکون مربوطه کلیک نمایید
 - 2. انتخاب اسلاید جهت آپلود تصویر
- 3. روی اسلاید مورد نظر کلیک کرده گزینه Save را انتخاب نمایید


Schwarz, Kathleen B., Schwarz, Kathleen B.... Show all. © 2018.

Fig. 65-1 (Adapted from http://tpis.upmc.com/tpislibrary/HB/H00034f.html .) Adenovirus hepatitis showing characteristic intranuclear inclusion bodies. A high magnification of the adenoviral lesion shows the light microscopic appearance of the typical adenoviral nuclear inclusions. The "smudgy" appearance of the nucleus (arrows), the peripheralized chromatin pattern, and the lack of cytoplasmic inclusions that help differentiate these inclusions from those seen with CMV.


برای ایجاد اسلاید جدید۱-گزینه Create New را کلیک نمایید. تا پنجره جدید گشوده شود ۲- سپس برای اسلاید خود نامی متناسب داده کلید Save را کلیک نمایید ۳- در مرحله بعد جهت افزودن تصویر گزینه Add and view presentation را کلیک کنید.




مشاهده تصاوير اضافه شده



cardiac Change











+ Add images from Saved Content







نحوه خروجی گرفتن از پاورپوینت ایجاد شده در پایگاه Clinical Key

حذف تصویر اضافه شده به اسلاید


Presentation Maker

My Presentations

+ Create new presentation

cardiac 10/8/17

cardiac Change







Export 13

IMAGE


Setup for cardiac surgery

Cardiac Surgery: Operative Technique.

Doty, Donald B., M.D.; Doty, John R., M.D... Published January 2, 2012.


A Operations are performed during a single aortic occlusion

ناريخچه جستجو



Drug Monographs

Browse Drug Monographs



۱- عنوان دارو را در باکس مربوطه تایپ کنید ۲- نام داروی مورد نظر را از لیست الفبایی انتخاب نمایید

اگر روی عنوان دارو کلیک نمایید اطلاعات کامل دارو را بدست می آورید. (مثال: صفحه اطلاعات داروی استامینوفن)

Indications & Dosage

Classifications

References

Global Drug Names


Acetaminophen; Clemastine; Pseudoephedrine

Tavist Allergy Sinus Headache

Drug Information Provided By Gold Standard


NOTE: In January 2007, the CDC warned caregivers and healthcare providers of the risk for serious injury or fatal overdose from the administration of cough and cold products to children and infants less than 2 years of age. 33534 This warning followed an investigation of the deaths of three (3) infants less than 6 months of age that were attributed to the inadvertent inappropriate use of these products. The symptoms preceding these deaths have not been clearly defined, and there is a lack of conclusive data describing the exact cause of death. The report estimated that 1519 children less than 2 years of age were treated in emergency departments during 2004-2005 for adverse events related to cough and cold medications, In October 2007, the FDA Nonprescription Drug Advisory Committee and the Pediatric Advisory Committee recommended that nonprescription cough and cold products containing pseudoephedrine, dextromethorphan, chlorpheniramine, diphenhydramine, brompheniramine, phenylephrine, clemastine, or guaifenesin not be used in children less than 6 years of age. In January 2008, the FDA issued a Public Health Advisory recommending that OTC cough and cold products not be used in infants and children less than 2 years. An official ruling regarding the use of these products in children greater than 2 years has not yet been announced. The FDA recommends that if parents and caregivers use cough and cold products in children greater than 2 years, labels should be read carefully, caution should be used when administering multiple products, and only measuring devices specifically designed for use with medications should be used. While some combination cough/cold products containing these ingredients are available by prescription only and are not necessarily under scrutiny by the FDA, clinicians should thoroughly assess each patient's use of similar products, both prescription and nonprescription, to avoid duplication of therapy and the potential for inadvertent overdose.

Clinical Trials



Guidelines

Browse Guidelines



Filter List by Title # A B C D E F G H I J K L M N O P Q R S T U V W X Y Z کلیدواژه مورد نظر خود را تایپ کنید و یا از لیست الفبایی انتخاب نمایید

(1) American Society of Clinical Oncology clinical practice guideline: update on adjuvant enc with hormone receptor-positive l American Society of Clinical Oncology - Medical Specialty So (1) Gonococcal infections. In: Sex treatment guideline 2010. (2) U Evidence Supporting the Recommendations transmitted diseases treatment gu Benefits/Harms of Centers for Disease Control and Prevention - Federal Gover Recommendations

(1) Peginterferon alfa and ribaviri Qualifying Statements chronic hepatitis C. (2) Peginterfe treatment of chronic hepatitis C. 2 National Institute for Health and Care Excellence (NICE) - No. National Healthcare Quality Report Categories Identifying Information and

Availability

Disclaimer

GUIDELINE

(1) Gonococcal infections. In: Sexually transmitted diseases treatment guidelines, 2010. (2) Update to CDC's sexually transmitted diseases treatment guidelines, 2010: oral cephalosporins no longer a recommended treatment for gonococcal infections.

n Rep. 2011 Jan 14:60(1):181, MMWR Recomm Rep. 2010 Dec 17:59(RR-12):49-55.

Recommendations

Major Recommendations


خلاصه ای از هر موضوع یا بیماری

Note from the National Guideline Clearinghouse (NGC) and the Centers for Disease Control and Prevention (CDC): On August 10, 2012, the CDC released an addendum to their 2010 recommendation for treatment of gonococcal infections. This report, which uses data from CDC's Gonococcal Isolate Surveillance Project (GISP), describes laboratory evidence of declining cefixime susceptibility among urethral Neisseria gonorrhoeae isolates collected in the United States during 2006-2011. Based on this data, CDC no longer recommends cefixime at any dose as a first-line regimen for treatment of gonococcal infections. The updated recommendations are presented below, followed by the recommendations from the original 2010 guideline.

2012 Addendum

Patient Education

Browse Patient Education





2,3-Diphosphoglycerate Test

24-Hour Urine Collection


5'-Nucleotidase Test

Available to print in English & Spanish. ExitCare, LLC. May 22, 2017.

Available to print in English & Spanish, ExitCare, LLC, May 22, 2017.

Available to print in English, Arabic... Show all, ExitCare, LLC, May 22, 2017.


کلیدواژه مورد نظر خود را تایپ کنید و یا از لیست الفبایی انتخاب نمایید



Nausea/Vomiting


First Consult/Clinical Overviews

Browse First Consult/Clinical Overviews



دسترسی سریع به آخرین و معتبرترین اطلاعات پزشکی مبتنی بر شواهد جهت ارزیابی، تشخیص، درمان و مدیریت بیماری است # A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

Browse First Consult/Clinical Overviews



Key points

The development of allergy to latex is asymptomatic, but once established, subsequent exposure to latex allergen can result in dermatitis (most often of the hands upon exposure to latex gloves), urticaria, conjunctivitis,

- While the prevalence of latex allergy in the general population is low, the risk of latex sensitization is higher in persons with regular x exposure
- There are no long many settings in which the use of latex is high; however, settings in which the use of powdered of gloves continues allow for significant potential exposure
- oince the late 1990s, the U.S. Food and Drug Administration has required manufacturers to apply warning labels to all products containing natural rubber latex (NRL), which eliminates the occult latex exposure that had contributed to sensitization and to unintentional exposure of sensitized individuals
- Primary prevention of latex sensitization and secondary prevention of latex allergy—related symptoms can be accomplished by reducing latex exposure for persons at risk
- Immediate hypersensitivity to latex is most reliably and safely confirmed with a consistent history of allergy symptoms after latex allergen exposure and with latex-specific serum immunoglobulin E (IgE) measurement

Background Description


FIRST CONSULT

Latex allergy

Revised: December 18, 2012

rhinitis, asthma, and anaphylaxis

Procedure consult



Sort by: Relevance >

[+] Rate Results

فراً یندهای پزشکی درمان بیماری را به صورت مرحله به مرحله همراه با فیلم ارائه می کند.



Video 1 - Topical amiodarone dur surgery: Does epicardial applicati event...

amiodarone-hydrogel, 150 e (3 mL) is added to a 2 ml amiodarone hydrock ximately 1.5 mL of amiod

each of the 2 syringes of the COSEAL kit. The mixe

ptember 1, 2017. Volu Greenstein, David, BM; Beau, Jordan, BSc... Show all. Publish

Catheter Ablation of Atrial Fibrillation (Cardiology

Intraoperative Transesophageal Echocardiography

VIDEO

Video 1 - Topical amiodarone during cardiac surgery: Does epicardial application of amiodarone prevent postoperative atrial fibrillation?

Journal of Thoracic and Cardiovascular Surgery, The

Greenstein, David, BM; Beau, Jordan, BSc... Show all.. Published September 1, 2017. Volume 154, Issue 3. Pages 886-892. © 2017.

For patients treated with amiodarone-hydrogel, 150 mg of liquid amiodarone hydrochloride (3 mL) is added to a 2 mL COSEAL kit (Baxter Healthcare). Approximately 1.5 mL of amiodarone is added to each of the 2 syringes of the COSEAL kit. The mixed a...



معاونت تحقيقات وفناوري

از توجه شما سپاسگزاریم

پست الكترونيكي جهت تماس

s.golmaghani@arums.ac.ir